

NPS55-78-033

NAVAL POSTGRADUATE SCHOOL  
Monterey, California



SIMPLE MODELS FOR POSITIVE-VALUED  
AND DISCRETE-VALUED TIME SERIES  
WITH ARMA CORRELATION STRUCTURE

by

P. A. W. Lewis

November 1978

Approved for public release; distribution unlimited.

FEDDOCS  
D 208.14/2:  
NPS-55-78-033

NAVAL POSTGRADUATE SCHOOL  
Monterey, California

Rear Admiral T. F. Dedman  
Superintendent

J. R. Borsting  
Provost

Reproduction of all or part of this report is authorized.

This report was prepared by:

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NPS55-78-033	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Simple Models for Positive-Valued and Discrete-Valued Time Series with ARMA Correlation Structure		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) P. A. W. Lewis		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, Ca. 93940		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61153N, RR014-05-01 N6001478WR80035
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		12. REPORT DATE November 1978
		13. NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Models, Discrete-Valued Time Series Positive-Valued Time Series ARMA correlation ARMA processes	Point processes Marginal distribution EARMA-type processes DARMA-type processes Autoregressive processes	Moving average processes
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Three models for positive-valued and discrete-valued stationary time series are discussed. All have the property that for a range of specified marginal distributions the time series have the same correlation structure as the usual linear, autoregressive-moving average (ARMA) model. The models differ in the range of marginal distributions which can be accommodated and in the simplicity and flexibility of each model. Specifically the EARMA-type processes can be extended from the exponential distribution to a rather narrow		

Continued

## 20. Abstract continued

range of continuous distributions; the DARMA-type processes can be defined usefully for any discrete marginal distribution and are simple and flexible. Finally the marginally controlled semiMarkov generated process can be defined for any continuous or discrete positive-valued distribution and is therefore very flexible. However the model suffers from some complexity and parametric obscurity.

SIMPLE MODELS FOR POSITIVE-VALUED AND DISCRETE-VALUED TIME SERIES  
WITH ARMA CORRELATION STRUCTURE

P. A. W. Lewis<sup>\*</sup>

Department of Operations Research  
Naval Postgraduate School  
Monterey, CA 93940

<sup>\*</sup> Paper presented at the Fifth International Symposium on Multivariate Analysis.  
To appear in the Proceedings published by North-Holland: Amsterdam.



Simple Models for Positive-Valued and Discrete-Valued  
Time Series with ARMA Correlation Structure

P. A. W. Lewis<sup>\*</sup>

Department of Operations Research  
Naval Postgraduate School  
Monterey, California 93940

Abstract

Three models for positive-valued and discrete-valued stationary time series are discussed. All have the property that for a range of specified marginal distributions the time series have the same correlation structure as the usual linear, autoregressive-moving average (ARMA) model. The models differ in the range of marginal distributions which can be accommodated and in the simplicity and flexibility of each model. Specifically the EARMA-type processes can be extended from the exponential distribution to a rather narrow range of continuous distributions; the DARMA-type processes can be defined usefully for any discrete marginal distribution and are simple and flexible. Finally the marginally controlled semi-Markov generated process can be defined for any continuous or discrete positive-valued distribution and is therefore very flexible. However the model suffers from some complexity and parametric obscurity.

\* Research supported by National Science Foundation Grant AF476 and Office of Naval Research Grant NR-42-284 at the Naval Postgraduate School.





## 1. Introduction

In much of the current work on the analysis of stationary time series there is an implicit assumption that the marginal distribution of the time series is normal. The assumption is implicit in that the marginal distribution is not considered to be of interest per se in the analysis, and also in that the statistical procedures which are used are very definitely based on normality assumptions. The stationary model on which much of this time series analysis is based is the mixed autoregressive moving average process,

$$a_0 X_i + a_1 X_{i-1} + \dots + a_p X_{i-p} = b_0 \epsilon_i + b_1 \epsilon_i + \dots + b_q \epsilon_{i-q} \quad (1.1)$$

$$i=0, \pm 1, \pm 2, \dots,$$

sometimes called the ARMA(p,q) or Box-Jenkins process. The process (1.1) is specified quite generally as a linear combination of i.i.d. random variables  $\{\epsilon_i\}$  of unspecified distribution, the linear, additive structure determining the correlation structure of the stationary sequence  $\{X_i\}$  under well-known restrictions on the parameters. If one wants  $\{X_i\}$  to be a time series with normally distributed marginal distribution, this can be accomplished by taking the  $\epsilon_i$ 's to be normally distributed. The model is then completely specified.

There are, however, many situations in which observations occur serially and in which the marginal distribution is patently non-normal. For example, data on the number of occurrences of all known diseases in each week is kept by the National Center for Health Statistics. The data is not only discrete count data, but for many diseases it is mostly on the order of 0, 1, 2, 3, and very seldom above this.

It has been suggested that such non-normal data be handled by data transformations and this is probably appropriate if the data is only slightly non-normal. In other cases it seems reasonable to start afresh and develop models from scratch. In this paper we summarize attempts to do this for stationary time series which are known to be non-normal because of either positivity or discreteness or both. The essence of the models is that the marginal distribution is specified, as well as the correlation structure. More generally the models are required to be simple and flexible in the following senses:

- a) The models should be specified in terms of easily observed and measured quantifiers. When the models are stationary, these quantifiers would typically be
  - i) the marginal distribution, and
  - ii) second-order moments (correlations).
- b) The models should be parametrically parsimonious and hopefully parametrically meaningful.
- c) The models should be easy to generate on computers, i.e., they should be structurally simple; in fact it might be preferable for the models to have linear structure.
- d) The models should be easy to fit to data, both informally and formally.

The model (1.1) certainly has most of the above features, but it is not known in general how to specify the distribution of  $\epsilon_i$  so as to produce a given, continuous marginal distribution for the  $X_i$ 's. Moreover, it is clearly not possible to do this at all if the  $X_i$ 's are discrete random variables.

The work described in this paper on non-normal time series is joint work with D. P. Gaver, P. A. Jacobs and A. J. Lawrance. Although the work has much broader connotation, it will be described in the context in which it arose, that of the description of stochastic point processes, or series of events occurring in time. One way in which these point processes can be described is as a sequence of intervals between events  $\{X_i\}$ , which are of course positive-valued random variables. In the common case of a Poisson point process the  $X_i$ 's have an exponential distribution. However, as in the case of epidemics, point processes are generally observed as counts of events in successive fixed intervals and these are non-negative discrete valued random variables. For the Poisson process these counts are independent and Poisson distributed and this serves as the null model in the analysis of count data from point processes.

Three distinct models are discussed in the context of the analysis and description of point processes. All of them satisfy the requirements discussed above to some degree. The EARMA-type process described first has recently been extended to have a complete ARMA-type correlation structure, but the process cannot be extended to all continuous marginal distributions. Marginally controlled semi-Markov generated processes, on the other hand, give a complete analog to (1.1), but they do not have linear structure. They can also be extended to give processes with discrete marginal distributions. A simpler, random linear structure has been derived, however, which gives discrete processes with ARMA structure. These are DARMA-type processes and come closer than the other processes to fulfilling the requirements of simplicity and flexibility.

Further details on the processes are to be found in the references.

## 2. Interval Models: Sequences of continuous positive-valued random variables

Univariate point processes in continuous time can be described equally well through the structure of the intervals between events  $\{X_i\}$ , where the  $X_i$ 's are continuous and positive-valued random variables, or the counting process  $\{N(t)\}$ , where  $N(t)$  gives the number of events in  $(0, t]$  and is discrete and non-negative. We discuss the modelling of the intervals  $\{X_i\}$  first. Of course the applications of the models are much broader; the  $X_i$ 's might for instance be the magnitudes of successive shocks in a sequence of earthquakes or the successive response times of a computer to messages sent via a terminal.

### 2.1. The first-order autoregressive exponential model (EAR(1))

In a Poisson process the intervals  $\{X_i\}$  are independent and identically distributed (i.i.d.) random variables with exponential distribution

$$F_X(x) = 1 - e^{-\lambda x}, \quad \lambda > 0; \quad x \geq 0. \quad (2.1)$$

Several attempts have been made to generalize the Poisson process by making the  $X_i$  dependent, but with exponential or conditionally exponential marginal distributions (Cox, 1955). The simplest and only really successful attempt in the sense of broad applicability (Gaver and Lewis, 1978) gives a process called the EAR(1) model, derived from the following consideration.

A first-order autoregressive stochastic sequence is defined by the stochastic difference equation (a special case of (1.1))

$$X_i = \rho X_{i-1} + \epsilon_i, \quad i=0, \pm 1, \pm 2, \dots; \quad |\rho| < 1, \quad (2.2)$$

where the  $\epsilon_i$  are assumed to be an i.i.d. stationary random sequence.

If the  $\epsilon_i$  are normally distributed, so are the  $X_i$ . What must the distribution of the  $\epsilon_i$  be in order for the  $X_i$  sequence to be stationary with an exponential( $\lambda$ ) distribution? The answer is surprisingly easy (Gaver and Lewis, 1978).

Let  $0 \leq \rho < 1$ , and let  $\{E_i\}$  be an i.i.d. exponential( $\lambda$ ) sequence. Now let  $\epsilon_i$  be equal to zero with probability  $\rho$  and equal to  $E_i$  with probability  $1-\rho$ . Then we have

$$X_i = \begin{cases} \rho X_{i-1} & \text{probability } \rho, \\ \rho X_{i-1} + E_i & \text{probability } (1-\rho), \end{cases} \quad (2.3)$$

$$= \rho X_{i-1} + V_i E_i, \quad (2.4)$$

where  $\{V_i\}$  is an i.i.d. binary sequence and  $P\{V_i=0\} = 1 - P\{V_i=1\} = \rho$ . Moreover if we let  $X_0 = E_0$ , and define  $X_i$  as in (2.3), the resulting sequence is stationary for  $i=0,1, \dots$ .

The point process with the interval structure (4.3) is called the EAR(1) point process. It is a tractable model, and most of its important properties are given in Gaver and Lewis (1978). In particular we have that  $\rho(k) = \rho^k$ . This model is in a sense degenerate because it contains runs of  $X_i$  in which values are exactly  $\rho$  times the previous value; it could, however, be a reasonable model for point processes observed in computer systems (e.g., inter-arrival times of requests to a storage subsystem) in which the intervals have exponential marginal distributions but are dependent. Note that as defined the model can only

provide sequences  $\{X_i\}$  with positive serial correlations. We can, however, define the process to include negative correlations (Gaver and Lewis, 1978); there is also a way to obviate the degeneracy (Lawrance, 1978).

Simple generalizations of this first-order, autoregressive, Markovian exponential process are the following.

## 2.2. The moving average exponential model (EMA(q)).

We define another stationary sequence  $\{X_i\}$ , using the  $\{E_i\}$  sequence above, according to

$$X_0 = E_0 , \quad (2.5)$$

$$X_i = \beta E_i + U_i E_{i-1}, \quad i=1, \dots; \quad 0 \leq \beta \leq 1 , \quad (2.6)$$

where  $\{U_i\}$  is an i.i.d. binary sequence in which  $U_i = 1$  with probability  $(1-\beta)$ . This is a first order exponential moving average process (EMA(1)) (Lawrance and Lewis, 1977) which is one-dependent; in particular

$$\rho(1) = \beta(1-\beta) \quad (2.7)$$

$$\rho(k) = 0 , \quad k=2, 3, \dots . \quad (2.8)$$

Properties of the EMA(1) process are given by Lawrance and Lewis (1977).

It is easy to see that we can make  $E_{i-1}$  in (2.6) a random linear combination of  $E_{i-1}$  and  $E_{i-2}$  to get an EMA(2) process, and can continue the process back  $q$  steps to obtain an EMA( $q$ ) process. The general EMA( $q$ ) model takes the form



$$X_i = \begin{cases} \beta_q E_i & \text{w.p. } b_{q+1} , \\ \beta_q E_i + \beta_{q-1} E_{i-1} & \text{w.p. } b_q , \\ \dots\dots\dots & \dots\dots\dots \\ \beta_q E_i + \beta_{q-1} E_{i-1} + \dots + \beta_1 E_{i-q+1} & \text{w.p. } b_2 , \\ \beta_q E_i + \beta_{q-1} E_{i-1} + \dots + \beta_1 E_{i-q+1} + E_{i-q} & \text{w.p. } b_1 , \end{cases} \quad (2.9)$$

for  $0 \leq \beta_1, \beta_2, \dots, \beta_q \leq 1$ ;  $i=0, \pm 1, \pm 2, \dots$  and

$$b_i = \begin{cases} \beta_q & i = q+1 , \\ (1-\beta_q) \dots (1-\beta_i) \beta_{i-1} & q \geq i \geq 2 , \\ (1-\beta_q) \dots (1-\beta_i) & i = 1 . \end{cases} \quad (2.10)$$

Note that the  $\beta_i$ 's can be obtained uniquely from the  $b_i$ 's; there are  $q+1$   $b_i$ 's but only  $q$   $\beta$ 's, since the sum of the  $b_i$ 's is equal to one.

This model is clearly only  $q$  dependent; in particular the correlations for the EMA( $q$ ) process are

$$\rho^{(q)}(k) = \text{corr}(X_i, X_{i-k}) = \begin{cases} \sum_{v=1}^{q-k+1} b_v b_{v+k} & 1 \leq k \leq q , \\ 0 & q+1 \leq k < \infty . \end{cases} \quad (2.11)$$

Thus the serial correlations are just lagged products of the  $b_i$  sequence and the formula (2.11) is completely analogous to the formula for the serial correlations of the standard MA( $q$ ) process; see Box and Jenkins (1970, p. 68). It can be seen from (2.11) that all the correlations are nonnegative and it may be further shown that they are bounded above by  $1/4$ .

### 2.3. The EARMA(1,1) model.

By making  $E_{i-q}$  in (2.9) autoregressive over the previous  $E_i$ 's, we obtain a mixed  $q$ th order moving-average, first order autoregressive process which we denote by EARMA(1, $q$ ). Consider explicitly the case  $q=1$ . The first order moving-average and first order autoregressive process EARMA(1,1) is given by

$$X_i = \beta E_i + U_i A_{i-1} , \quad (2.12)$$

with

$$A_{i-1} = \rho A_{i-2} + V_i E_{i-1} , \quad (2.13)$$

for  $i=1, 2, 3, \dots$  and  $A_{-1} = E_{-1}$  with  $U_i$  and  $V_i$  as defined above. This sequence of random variables is not Markovian.

The second-order correlation structure of the process is given by

$$\rho(k) = \rho^{k-1} c(\beta, \rho) , \quad (2.14)$$

where

$$c(\beta, \rho) = \beta(1-\beta) + \rho(1-\beta)(1-2\beta) . \quad (2.15)$$

The point process whose intervals have the EARMA(1,1) structure is discussed in detail in Jacobs and Lewis (1977). In particular, for  $\beta=1$  it is a Poisson process. The process is very simple to generate on a computer and is very useful for modelling dependent sequences in queueing systems (Jacobs, 1978; Lewis and Shedler, 1978).

### 2.4. The $p$ th-order autoregressive model EAR( $p$ ).

Quite recently ways have been found to obtain exponential sequences  $\{X_i\}$  which have autoregressive structure of order  $p$ , and to combine these with the moving average process to get a mixed autoregressive-moving



average process EARMA(p,q); see Lewis and Lawrance (1978). Another method of defining pth-order autoregressive exponential sequences, which is closely related to the DARMA(p,q) process discussed later, and which we have only just begun to explore, is described here.

This pth-order exponential autoregressive model can be written as

$$X_i = \alpha_{S_i} X_{i-S_i} + \epsilon_{i,S_i}, \quad (2.16)$$

where the  $S_i$ 's are i.i.d. discrete random variables taking values 1, 2, ..., p, and  $\epsilon_{i,S_i}$  is defined to be 0 w.p.  $\alpha_j$ , and  $E_i$  w.p.  $\alpha_j$  if  $S_i = j$ . If one assume stationarity and that  $X_{i-1}, X_{i-2}, \dots$  are marginally exponential( $\lambda$ ), then  $X_i$  is a random mixture of  $E_i$  and  $X_{i-1}, \dots, X_{i-p}$  and is exponential( $\lambda$ ). The correlation equations from this process are variants of the familiar Yule-Walker equations. The model is more tractable than the pth-order autoregressive process given in Lewis and Lawrance (1978) and is probably simpler to extend to other distributions than the exponential.

A drawback of these EARMA-type processes is that the serial correlations are all positive, although the scheme given in Gaver and Lewis (1978) for a negatively correlated EARMA1 process can probably be extended to the complete EARMA(p,q) process.

## 2.5. The semi-Markov generated point process with fixed marginal distribution.

The question arises as to whether there are interval processes  $\{X_i\}$  with exponential marginal distributions and, for example, ARMA(1,1) second-order correlation structure and which cover a broader range of correlation than the EARMA(1,1) process (though perhaps at a cost of more complicated structure).

We discuss briefly one such process. It is a special case of the semi-Markov generated point process introduced by Cox (1962) and extended by Haskell and Lewis (1978). We first describe the two-state semi-Markov generated model. In this model there are two types of intervals with distributions  $F_1(x)$  and  $F_2(x)$ , sampled in accordance with a two-state Markov chain for which the one-step transition matrix is

$$\underline{P} = \begin{pmatrix} \alpha_1 & 1-\alpha_1 \\ 1-\alpha_2 & \alpha_2 \end{pmatrix} \quad (2.18)$$

and the stationary vector is

$$\underline{\Pi} = \underline{\Pi} \underline{P} = \left( \frac{1 - \alpha_2}{2 - \alpha_1 - \alpha_2}, \frac{1 - \alpha_1}{2 - \alpha_1 - \alpha_1} \right). \quad (2.19)$$

When we form the point process we assume that no information is available about the type of interval, i.e., that in the actual bivariate point process of transitions we suppress knowledge of the type of transition. Then the distribution of an interval between transitions (events)  $X_i$  in the stationary point process is

$$F_X(x) = \pi_1 F_1(x) + \pi_2 F_2(x) \quad (2.30)$$

and the correlation between  $X_i$  and  $X_{i+k}$  is

$$\rho(k) = M^k, \quad k=1, 2, \dots, \quad (2.21)$$

where  $M$  is a positive constant and  $\beta = \alpha_1 + \alpha_2 - 1 = \alpha_1 (1 - \alpha_2)$ .

Thus the correlation structure is that of an ARMA(1,1) process. For a derivation of this result see Cox and Lewis (1966, Ch. 7, 194-196).

Lewis and Shedler (1973) used this process to model the page exception process in a multiprogrammed computer system. The problem is to deal with the mixture distribution (2.20) for the marginal distribution of intervals; this seems to limit the utility of the model. However, there is a way around it which produces a marginally controlled semi-Markov generated process.

To obtain an exponential marginal distribution, consider the following device (Jacobs and Lewis, 1977). Fix  $x_0$ , where  $0 < x_0 < \infty$ , and let

$$F_1(x) = \begin{cases} \frac{\int_0^x e^{-\lambda u} du}{1 - e^{-\lambda x_0}} & 0 \leq x \leq x_0, \\ 1 & x > x_0; \end{cases} \quad (2.22)$$

$$F_2(x) = \begin{cases} 0 & x \leq x_0, \\ \frac{\int_0^x e^{-\lambda u} du}{e^{-\lambda x_0}} & x > x_0; \end{cases}$$

then  $F(x)$ , the marginal distribution of an interval, is exponential( $\lambda$ ) if we set  $\pi_1 = 1 - \exp(-\lambda x_0)$ . There is one degree of freedom left in the matrix  $\underline{P}$ ; in addition to  $\lambda$ , we have free parameters  $\pi_1$  (or  $x_0$ ) and  $\alpha_1$  although the range of  $\alpha_1$  is restricted. What then is the range of  $\beta$ , and can it be negative?

Straightforward manipulation shows that

$$\beta = \frac{\pi_1 - \alpha_1}{\pi_1 - 1}, \quad (2.23)$$

which lies in absolute value between zero and one but can be negative; therefore the serial correlations can be negative. Thus the model appears to be broader than the EARMA(1,1) model. The question of comparing the two models when  $\beta$  is positive has not yet been explored; it requires higher order interval correlations, as discussed by Brillinger (1972).

## 2.6. Generalizations

The marginally controlled semi-Markov generated sequence  $\{X_i\}$  discussed above can be extended in such a way that  $X_i$  will have any distribution, say  $F(x)$ . Thus we let

$$\begin{aligned} F_1(x) &= \begin{cases} \frac{F(x)}{F(x_0)} & 0 \leq x \leq x_0, \\ 1 & x > x_0; \end{cases} \\ F_2(x) &= \begin{cases} 0 & x \leq x_0, \\ \frac{F(x) - F(x_0)}{1 - F(x_0)} & x > x_0; \end{cases} \end{aligned} \quad (2.24)$$

then the marginal distribution of an interval is equal to  $F(x)$ , from (2.30), if we set  $\pi_1 = F(x_0)$ . Note that the model is very non-linear and the correlation structure is a complicated function of the functional form of  $F(x)$ .

The much simpler EARMA structure can be extended to some extent. Random variables for which the equation (2.2) has a proper solution are called self-decomposable random variables on random variables of type L. This class includes random variables with Gamma, Cauchy, Pareto, double exponential and perhaps many other distributions. For these random

variables, a  $p$ th-order-autoregressive process can be defined as at (2.16). The unique feature of the exponential process is that the  $\epsilon_i$  which makes  $X_i$  exponential( $\lambda$ ) in (2.2) is again an exponential( $\lambda$ ) random variable, albeit mixed with an atom at zero. This property, shared with the double exponential and normal random variables, is what makes it simple to define a moving-average type process, as at (2.9).

### 3. Count Models: Sequences of discrete-valued random variables.

As remarked earlier, most data on point processes is recorded as numbers of events in successive fixed-length intervals. Despite this fact, most point process models assume that exact times of events are known and it is not simple to derive from these models the statistics of the counts in fixed intervals. Thus in this area in particular flexible models for discrete-valued random variables are needed.

Another application might be to modelling of air pollution data in which concentrations of various chemicals in the air is indicated on a scale of zero to ten. In general this situation requires multivariate time series, but space prohibits discussion of multivariate versions of the DARMA-type processes discussed in this section.

#### 3.1. The first-order autoregressive discrete model (DAR(1)).

Again we denote the sequence of discrete-valued random variables by  $\{X_i\}$ . If the  $X_i$  are counts in a Poisson process then the  $X_i$ 's are i.i.d. Poisson-distributed random variables. Once dependence is observed in data it is useful to assume, as a first cut, that the dependence is Markovian and use a Markov chain model in which the distribution of  $X_{i+1}$  depends only on the value of  $X_i$  and is specified by the transition matrix  $\underline{P}$  with elements

$$P(k,j) = P\{X_{i+1} = j | X_i = k\} , \quad (3.1)$$

with  $j$  and  $k$  taking values in the space  $E$ , a discrete subset of the real line. Under suitable conditions there is a stationary distribution  $\pi$  for  $\{X_i\}$  given by the equation

$$\pi = \pi P . \quad (3.2)$$

The Markov chain model (3.2) is by virtue of its place in the statistician's toolbox the discrete counterpart of the AR1 process. However the AR1 process has one dependency parameter  $\rho$ , plus any parameters which specify the distribution of the  $\epsilon_i$ 's. The Markov chain on the other hand can have an infinite number of parameters and in many cases  $\pi$  cannot be obtained explicitly from (3.2). This is awkward for statistical analysis. A solution is given by constructing the DAR(1) model (discrete autoregressive model of under one) which is an analog of the EAR(1) model, as follows.

Let  $Y_i$  be an i.i.d. sequence of random variables taking values in the space  $E$ , and let  $V_i$  be an i.i.d. binomial sequence for which  $P\{V_i = 1\} = \rho$ . Then

$$X_i = V_i X_{i-1} + (1 - V_i)Y_i \quad i=0, \pm 1, \pm 2, \dots; \quad 0 \leq \rho < 1 . \quad (3.3)$$

$$= \begin{cases} X_{i-1} & \text{w.p. } \rho , \\ Y_i & \text{w.p. } (1-\rho) . \end{cases} \quad (3.4)$$

If  $X_0$  has distribution  $\pi$ , then so does  $X_1$  since it is a mixture of two random variables,  $X_0$  and  $Y_1$ , with distribution  $\pi$ . Consequently all the  $X_i$ ,  $i=1, 2, \dots$  have marginal distribution  $\pi$ .



Note that  $\{X_i\}$  is a Markov chain with transition probabilities

$$P(k,j) = P\{X_{i+1} = j | X_i = k\} = \begin{cases} (1-\rho) \pi(j) & k \neq j, \\ \rho + (1-\rho) \pi(j) & k = j; \end{cases} \quad (3.5)$$

in fact it is a Markov chain in which the correlation structure is specified by one parameter  $\rho$ , and with specified marginal (stationary) distribution  $\underline{\pi}$ . Thus  $\pi$  may be a Poisson distribution and then the DAR1 model is a 2-parameter  $(\lambda, \rho)$  Markov chain. The analogy with the AR(1) model is clear

As with the EAR(1) model the serial correlations are  $\rho(k) = \rho^k \geq 0$ . Extensions to negatively correlated sequences are given in Jacobs and Lewis (1978).

### 3.2. The pth-order autoregressive discrete model (DAR(p)).

First order Markov dependence is a special kind of dependence which is attractive because of analytical tractability considerations, but it is not necessarily met with in practice. One immediate consequence of the Markovian property is that runs of distinct values, say  $X_i = j$ , have a length which is geometrically distributed (Jacobs and Lewis, 1978a) and this is easily checked in data. If the data fails to have this property, what other types of dependency can be utilized?

A first direction might be to go to higher order (say pth-order) autoregression, which is an explicit pth-order Markov structure, and the DAR(1) model can be extended in this direction. Thus in addition to the assumptions at (3.3) let  $A_i$  be an i.i.d. sequence of random variables taking values in  $\{1, 2, \dots, p\}$ , with  $P\{A_i = j\} = \alpha_j$ . Then the DAR(p) process is defined as

$$X_i = v_i X_{i-A_i} + (1 - v_i)Y_i, \quad i=0, \pm 1, \pm 2, \dots \quad (3.6)$$

so that  $X_i$  is (exclusively) either one of the previous  $p$  values  $X_{i-1}, \dots, X_{i-p}$ , or the error term  $Y_i$ . Properties of this model are developed extensively in Jacobs and Lewis (1978c). When  $\alpha_1 = 1$ , and all other  $\alpha_j$ 's are zero it is the DAR(1) model.

Yule-Walker equations for the correlations in the stationary DAR(p) process are given in Jacobs and Lewis (1973c) as well as stationarity conditions. In particular for  $p=2$  we have the limiting result

$$v(k,j) = \lim_{i \rightarrow \infty} P\{X_{i+1}=k, X_{i+2}=j\} = \begin{cases} \{1-\rho(1)\}\pi(k)\pi(j) & k \neq j, \\ \rho(1)\pi(j) + \{1-\rho(1)\}\pi(j)^2 & k = j, \end{cases} \quad (3.7)$$

where  $\rho(1) = \text{corr}(X_i, X_{i+1})$  in the stationary process. Thus, if we let  $X_0$  and  $X_{-1}$  have the joint distribution  $v(k,j)$ , a stationary, second-order autoregressive process with any marginal distribution can be generated. A scheme for obtaining sequences which are possibly negatively correlated is given in Jacobs and Lewis (1978c).

### 3.3. The q-th order moving average discrete model (DMA(q)).

The other alternative to Markovian dependence (of any order) which is usually considered in time series analysis is the finite-length dependence produced by the moving-average part of the ARMA(p,q) process (1.1). This type of behavior is easily produced for discrete random variables by a random index model of the type

$$X_i = Y_{i-S_i}, \quad (3.8)$$



where  $S_i$  are i.i.d. random variables with  $P\{S_i \leq k\} = b_k$ . Thus we may write

$$X_i = Y_{i-k} \quad \text{w.p.} \quad b_k - b_{k-1}, \quad k=0, \dots, q; \quad b_{-1} = 0. \quad (3.9)$$

The autoregressive process DAR(p) is also a random index model, but the random indices are not independent. The correlation structure of this DMA(q) process is easily found to be

$$\begin{aligned} \rho^{(q)}(k) = \text{corr}(X_i, X_{i-k}) &= \sum_{v=0}^{q-k} b_v b_{v+k} & 1 \leq k \leq q, \\ &= 0 & k > q. \end{aligned} \quad (3.10)$$

This is the exact analog of (2.11) for the EMA(q) process and the corresponding formula for the MA(q) process. Note that the DMA(q) process is not Markovian. Runs properties of the process are given in Jacobs and Lewis (1978a); the runs are not geometrically distributed.

#### 3.4. Mixed autoregressive-moving average discrete models.

As in the case of the ARMA(p,q) model (1.1), it is useful to have both autoregressive, Markovian dependence and moving average dependence combined into one model. In Jacobs and Lewis (1978a) this was done by replacing the  $Y_{i-q}$  term in (3.8) by a discrete autoregression (3.3) over  $Y_{i-q}, Y_{i-q-1}, \dots$ . Clearly this can be extended by replacing  $Y_{i-q}$  by a p-th order autoregression (3.6) over  $Y_{i-q}, Y_{i-q-1}, \dots$  to obtain a DARMA(p,q) model which is the analog of the EARMA(p,q) model of Lawrance and Lewis (1978). This is not a complete analog of the ARMA(p,q) model in that there is no cross-over of the autoregression and

the moving average, but it is in fact possible to do this to obtain a model called NDARMA(p,q) as follows:

Let

$$X_i = V_i X_{i-A_i} + (1 - V_i) Y_{i-S_i} \quad i=0, \pm 1, \pm 2, \dots, \quad (3.11)$$

where the  $A_i$  are i.i.d. random variables taking values in  $\{1, 2, \dots, p\}$  with  $P\{A_i=j\} = \alpha_j$ ; the  $S_i$  are i.i.d. random variables taking values in  $\{0, \dots, q\}$  with  $P\{S_i \leq k\} = F(k)$  and the  $V_i$ 's are i.i.d. Bernoulli random variables with  $P\{V_i=1\} = \rho$ .

The model works because a mixture of dependent random variables, all with marginal distribution  $\pi$ , has distribution  $\pi$ ; thus if  $X_{i-1}, \dots, X_{i-p}$  have marginal distribution  $\pi$ , then so will  $X_i$  since it is a mixture of the dependent random variables  $X_{i-1}, \dots, X_{i-p}$  and  $Y_i, \dots, Y_{i-q}$ . Note that when  $\rho=0$  we have the DMA(q) process; if in addition  $F(0) = 1$  the sequence is i.i.d. since  $X_i = Y_i$ . When  $1 > \rho \neq 0$ ,  $F(0) = 1$  we have the DAR(p) process. Thus the parameters are such that interesting special cases fall out easily. Moreover the  $\rho$  parameter measures the degree of mixture of Markovian and moving average dependence, and the distributions of the  $A_i$ 's and  $S_i$ 's give a picture of where the dependence is lagged over previous  $X_i$  or  $Y_i$  values.

The model (3.11) has not yet been fully explored. At first sight it seems preferable to the DARMA(p,q) model, possibly because of the compactness of (3.11) and its close analogy to ARMA(p,q) models. The DARMA(p,q) and NDARMA(p,q) models are, however, distinct and in fact preliminary investigation of the (1.1) case shows that the DARMA(1,1) model (Jacobs and Lewis, 1978a) has a broader correlation structure than does the

NDARMA(1,1). On the other hand the autoregression is not explicit in the DARMA(1,1) model. Both models, therefore, will probably be useful in modelling discrete data such as occur in sampled point processes.

### 3.5. The marginally controlled semi-Markov generated process.

In the structure of the 2-state marginally controlled semi-Markov generated process detailed at (2.24) no assumption was made about continuity of  $F(x)$ . Thus  $F(x)$  could be discrete, giving a sequence  $\{X_i\}$  with known discrete marginal distribution  $F(x)$  and ARMA(1,1) correlation structure. By going to an  $n$ -state semi-Markov model, a process with ARMA( $p,q$ ) correlation structure can be generated (Haskell and Lewis, 1978) with  $n$  a function of  $p$  and  $q$ , and the procedure to obtain a given marginal distribution is just an extension of (2.24). Thus we have, in terms of the quantification of the process by marginal distribution and correlation structure, a direct competitor to the DARMA-type processes.

Comparison of the two types of discrete processes is interesting and points up the simplicity of the DARMA-type processes. In particular the correlation structure of the DARMA( $p,q$ ) process is explicit in form if not in detail and the process is a simple, random linear combination of random variables generated from an i.i.d. sequence  $Y_i$ . This is clearly not true for the marginally controlled semi-Markov generated process; the recognition that its correlation structure is ARMA-type is accidental and not intuitive. Deeper comparison of these processes in terms, say, of the range of correlation the model will encompass will be instructive. Here again the DARMA-type processes have an advantage; their correlation structure is independent of the marginal distribution  $\pi$ .

#### 4. Summary and Conclusions

We have outlined in this paper three models for discrete-valued and positive-valued time series, all of which to some degree satisfy the criteria of flexibility or simplicity or both set forth in the introduction. Perhaps the main point about the models is that they are designed to accomodate situations in which the marginal distributions in the stationary processes are given and are non-normal.

Properties of these models such as mixing and asymptotic results, higher-order moments, distributions of runs for the discrete models and sums of random variables and point spectra are considered in the references.

There are many other properties of the processes which are still to be explored. Statistical estimation, except in an ad hoc manner and for the Markovian cases, is difficult and has yet to be examined. Extensions to multivariate cases is of great interest for real applications and has been done to some degree in the context of queues with correlated service and arrival times (Jacobs, 1978, and Lewis and Shedler, 1978). The DARMA-type processes, in particular, can be easily extended to coupled equations in the same way as linear processes are extended in econometric models. They might therefore find use in modelling multivariate situations such as the number of cars passing different points in a road evaluated in successive fixed time intervals.

Finally an important problem is to extend the models so as to include inhomogeneity, particularly of the seasonal type, and the effects of concomittant or auxilliary variables. Several schemes are under consideration for these extensions of the models.

## References

- Box, G. E. P. and Jenkins, G. (1970). Time Series Analysis, Forecasting and Control. San Francisco: Holden Day.
- Brillinger, D. R. (1972). The spectral analysis of stationary interval functions. Proc. 6th Berkeley Symp. Math. Stat. and Prob., Berkeley: University of California Press, 483-514.
- Cox, D. R. (1955). Some statistical methods connected with series of events. J. R. Statist. Soc. B, 17, 129-164.
- Cox, D. R. (1962). Renewal Theory. London: Methuen.
- Cox, D. R. and Lewis, P. A. W. (1966). The Statistical Analysis of Series of Events. London: Methuen and New York: Wiley.
- Gaver, D. P. and Lewis, P. A. W. (1978). First order autoregressive Gamma sequences. Naval Postgraduate School Report NPS5-78-016.
- Haskell, R. and Lewis, P. A. W. (1978). Interval spectra of semi-Markov generated point processes. To appear.
- Jacobs, P. A. (1978). A closed cyclic queueing network with dependent exponential service times. J. Appl. Prob., to appear.
- Jacobs, P. A. and Lewis, P. A. W. (1977). A mixed autoregressive-moving average exponential sequence and point process (EARMA(1,1)). Adv. Appl. Prob., 9, 87-104.
- Jacobs, P. A. and Lewis, P. A. W. (1978a). Discrete time series generated by mixtures. I: Correlational and runs properties. J. R. Statist. Soc. B, 40, 1, 94-105.
- Jacobs, P. A. and Lewis, P. A. W. (1978b). Discrete time series generated by mixtures. II: Asymptotic properties. J. R. Statist. Soc. B, 40, to appear.
- Jacobs, P. A. and Lewis, P. A. W. (1978c). Discrete time series generated by mixtures. III: Autoregressive processes (DAR(p)). To appear.
- Lawrance, A. J. (1978). Some autoregressive models for point processes. Proc. Bolyai Janos Math. Soc. Colloquium on Point Processes and Queueing Theory, to appear.
- Lawrance, A. J. and Lewis, P. A. W. (1977). A moving average exponential point process (EMA1). J. Appl. Prob., 14, 98-113.
- Lawrance, A. J. and Lewis, P. A. W. (1978). An exponential autoregressive-moving average process EARMA(p,q): Definition and correlational properties. Naval Postgraduate School Report NPS55-78-1.

- Lewis, P. A. W. and Shedler, G. S. (1973). Empirically derived micro-models for sequences of page exceptions. IBM J. Res. Dev., 17, 86-100.
- Lewis, P. A. W. and Shedler, G. S. (1978). Analysis and modelling of point processes in computer systems. Bull. Int. Statist. Inst., to appear.



# DISTRIBUTION LIST

NO. OF COPIES

STATISTICS AND PROBABILITY PROGRAM OFFICE OF NAVAL RESEARCH CODE 436 ARLINGTON VA	22217	1
OFFICE OF NAVAL RESEARCH NEW YORK AREA OFFICE 715 BROADWAY - 5TH FLOOR ATTN: DR. ROGER GRAFTON NEW YORK, NY	10003	1
DIRECTOR OFFICE OF NAVAL RESEARCH BRANCH OFF 536 SOUTH CLARK STREET ATTN: DEPUTY AND CHIEF SCIENTIST CHICAGO, IL	60605	1
LIBRARY NAVAL OCEAN SYSTEMS CENTER SAN DIEGO CA	92152	1
NAVY LIBRARY NATIONAL SPACE TECHNOLOGY LAB ATTN: NAVY LIBRARIAN BAY ST. LOUIS MS	39522	1
NAVAL ELECTRONIC SYSTEMS COMMAND NAVELEX 320 NATIONAL CENTER NO. 1 ARLINGTON VA	20360	1
DIRECTOR NAVAL RESEARCH LABORATORY ATTN: LIBRARY (ONRL) CODE 2029 WASHINGTON, D.C.	20375	1
DEFENSE DOCUMENTATION CENTER CAMERON STATION  ALEXANDRIA VIRGINIA 22314		1
TECHNICAL INFORMATION DIVISION NAVAL RESEARCH LABORATORY  WASHINGTON, D. C.	20375	1

	NO. OF COPIES
OFFICE OF NAVAL RESEARCH SAN FRANCISCO AREA OFFICE 760 MARKET STREET SAN FRANCISCO CALIFORNIA 94102	1
TECHNICAL LIBRARY NAVAL ORDNANCE STATION  INDIAN HEAD MARYLAND 20640	1
NAVAL SHIP ENGINEERING CENTER PHILADELPHIA DIVISION TECHNICAL LIBRARY PHILADELPHIA PENNSYLVANIA 19112	1
BUREAU OF NAVAL PERSONNEL DEPARTMENT OF THE NAVY TECHNICAL LIBRARY WASHINGTON D. C. 20370	1
LIBRARY CODE 0212 NAVAL POSTGRADUATE SCHOOL  MONTEREY CALIFORNIA 93940	1
PROF. M. AEDEL-HAMEED DEPARTMENT OF MATHEMATICS UNIVERSITY OF NORTH CAROLINA CHARLOTTE NC  28223	1
PROF. T. W. ANDERSON DEPARTMENT OF STATISTICS STANFORD UNIVERSITY  STANFORD , CALIFORNIA 94305	1
PROF. F. J. ANSCOMBE DEPARTMENT OF STATISTICS YALE UNIVERSITY NEW HAVEN CONNECTICUT 06520	1
PROF. L. A. ARCIAN INSTITUTE OF INDUSTRIAL ADMINISTRATION UNION COLLEGE SCHENECTADY , NEW YORK 12308	1
PROF. C. R. BAKER DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CAROLINA CHAPEL HILL, NORTH CAROLINA 27514	1



NO. OF COPIES

PROF. R. E. BECHOFER  
DEPARTMENT OF OPERATIONS RESEARCH  
CORNELL UNIVERSITY  
ITHACA  
NEW YORK 14850

1

PROF. N. J. BERSHAD  
SCHOOL OF ENGINEERING  
UNIVERSITY OF CALIFORNIA  
IRVINE,  
CALIFORNIA  
92664

1

P. J. BICKEL  
DEPARTMENT OF STATISTICS  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA

1

94720

PROF. F. W. BLICK  
DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF PITTSBURGH  
PITTSBURGH  
PA

1

15260

PROF. JOSEPH BLUM  
DEPT. OF MATHEMATICS, STATISTICS  
AND COMPUTER SCIENCE  
THE AMERICAN UNIVERSITY  
WASHINGTON  
DC

1

20016

PROF. R. A. BRADLEY  
DEPARTMENT OF STATISTICS  
FLORIDA STATE UNIVERSITY

1

TALLAHASSEE, FLORIDA 32306

PROF. R. E. BARLOW  
OPERATIONS RESEARCH CENTER  
COLLEGE OF ENGINEERING  
UNIVERSITY OF CALIFORNIA  
BERKELEY  
CALIFORNIA 94720

1

MR. C. N. BENNETT  
NAVAL COASTAL SYSTEMS LABORATORY  
CODE P761  
PANAMA CITY,  
FLORIDA  
32401

1

PROF. U. N. BIFAT  
COMPUTER SCIENCE / OPERATIONS  
RESEARCH CENTER  
SOUTHERN METHODIST UNIVERSITY  
DALLAS,  
TEXAS 75275

1

PROF. W. R. BLISCHKE  
DEPT. OF QUANTITATIVE  
BUSINESS ANALYSIS  
UNIVERSITY OF SOUTHERN CALIFORNIA  
LOS ANGELES, CALIFORNIA

1

## NO. OF COPIES

DR. DERRILL J. BORDELON  
NAVAL UNDERWATER SYSTEMS CENTER  
CODE 21  
NEWPORT  
RI

1

C2840

J. E. BOYER JR  
DEPT. OF STATISTICS  
SOUTHERN METHODIST UNIVERSITY  
DALLAS  
TX

1

75275

DR. J. CHANDRA  
U. S. ARMY RESEARCH  
P. O. BOX 12211  
RESEARCH TRIANGLE PARK,  
NORTH CAROLINA  
27706

1

PROF. F. CHERNOFF  
DEPT. OF MATHEMATICS  
MASS INSTITUTE OF TECHNOLOGY  
CAMBRIDGE,  
MASSACHUSETTS 02139

1

PROF. C. DERNAN  
DEPARTMENT OF CIVIL ENGINEERING  
AND ENGINEERING MECHANICS  
COLUMBIA UNIVERSITY  
NEW YORK  
NEW YORK

1

10027

PROF. R. L. DISNEY  
VIRGINIA POLYTECHNIC INSTITUTE  
AND STATE UNIVERSITY  
DEPT. OF INDUSTRIAL ENGINEERING  
AND OPERATIONS RESEARCH  
BLACKSBURG, VA

1

24061

MR. J. DOWLING  
DEFENSE LOGISTICS STUDIES  
INFORMATION EXCHANGE  
ARMY LOGISTICS MANAGEMENT CENTER  
FORT LEE,  
VIRGINIA 20390

1

PROF. J. D. ESARY  
DEPT. OF OPERATIONS RESEARCH AND  
ADMINISTRATIVE SCIENCES  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY,  
CALIFORNIA 93940

1

DR. M. J. FISCHER  
DEFENSE COMMUNICATIONS AGENCY  
1860 WIEHLE AVENUE  
RESTON  
VIRGINIA  
22070

1

PROF. C. P. GAVER  
DEPT. OF OPERATIONS RESEARCH  
NAVAL POSTGRADUATE SCHOOL  
MONTEREY  
CA

1

53940

MR. GENE F. GLEISSNER  
APPLIED MATHEMATICS LABORATORY  
DAVID TAYLOR NAVAL SHIP RESEARCH  
AND DEVELOPMENT CENTER  
BETHESDA  
MD

1

20084

PROF. S. S. GUPTA  
DEPARTMENT OF STATISTICS  
PURDUE UNIVERSITY  
LAFAYETTE  
INDIANA 47907

1

PROF. D. L. HANSON  
DEPT OF MATH. SCIENCES  
STATE UNIVERSITY OF NEW YORK,  
BINGHAMTON  
BINGHAMTON  
NY

1

13901

PROF. F. J. HARRIS  
DEPT. OF ELECTRICAL ENGINEERING  
SAN DIEGO STATE UNIVERSITY  
SAN DIEGO  
CA

1

52182

PROF. L. H. HERBACH  
DEPT. OF OPERATIONS RESEARCH AND  
SYSTEMS ANALYSIS  
POLYTECHNIC INSTITUTE OF NEW YORK  
BROOKLYN  
NY

1

11201

PROF. M. J. HINICH  
DEPARTMENT OF ECONOMICS  
VIRGINIA POLYTECHNIC INSTITUTE  
AND STATE UNIVERSITY  
BLACKSBURG,  
VIRGINIA 24061

1

PROF. W. M. HIRSCH  
INSTITUTE OF MATHEMATICAL SCIENCES  
NEW YORK UNIVERSITY  
NEW YORK  
NEW YORK 10453

1

PROF. C. L. IGLEHART  
DEPARTMENT OF OPERATIONS RESEARCH  
STANFORD UNIVERSITY  
STANFORD,  
CALIFORNIA  
94350

1

NO. OF COPIES

PROF. J. B. KACANE  
DEPARTMENT OF STATISTICS  
CARNEGIE-MELLON  
PITTSBURGH,  
PENNSYLVANIA  
15213

1

DR. RICHARD LAU  
DIRECTOR  
OFFICE OF NAVAL RESEARCH BRANCH OFF  
1030 EAST GREEN STREET  
PASADENA  
CA

1

91101

DR. A. R. LAUFER  
DIRECTOR  
OFFICE OF NAVAL RESEARCH BRANCH OFF  
1030 EAST GREEN STREET  
PASADENA  
CA

1

91101

PROF. M. LEADBETTER  
DEPARTMENT OF STATISTICS  
UNIVERSITY OF NORTH CAROLINA  
CHAPEL HILL  
NORTH CAROLINA 27514

1

DR. J. S. LEE  
J. S. LEE ASSOCIATES, INC.  
2001 JEFFERSON DAVIS HIGHWAY  
SUITE 8C2  
ARLINGTON  
VA

1

22202

PROF. L. C. LEE  
DEPARTMENT OF STATISTICS  
VIRGINIA POLYTECHNIC INSTITUTE  
AND STATE UNIVERSITY  
BLACKSBURG  
VA

1

24061

PROF. R. S. LEVENWORTH  
DEPT. OF INDUSTRIAL AND SYSTEMS  
ENGINEERING  
UNIVERSITY OF FLORIDA  
GAINSVILLE,  
FLORIDA 32611

1

PROF. G. LIEBERMAN  
STANFORD UNIVERSITY  
DEPARTMENT OF OPERATIONS RESEARCH  
STANFORD CALIFORNIA 94305

1

NO. OF COPIES

DR. JAMES F. MAAR  
NATIONAL SECURITY AGENCY  
FCPT MEADE , MARYLAND  
20755 1

PROF. R. W. MADSEN  
DEPARTMENT OF STATISTICS  
UNIVERSITY OF MISSOURI  
COLUMBIA  
MC 1

65201

DR. N. R. MANN  
SCIENCE CENTER  
ROCKWELL INTERNATIONAL CORPORATION  
P.O. BOX 1085  
THOUSAND OAKS ,  
CALIFORNIA 91360 1

DR. W. H. MARLOW  
PROGRAM IN LOGISTICS  
THE GEORGE WASHINGTON UNIVERSITY  
707 22ND STREET , N. W.  
WASHINGTON , D. C.  
20037 1

PROF. E. MASRY  
DEPT. APPLIED PHYSICS AND  
INFORMATION SERVICE  
UNIVERSITY OF CALIFORNIA  
LA JOLLA  
CALIFORNIA 92093 1

DR. BRUCE J. McDONALD  
SCIENTIFIC DIRECTOR  
SCIENTIFIC LIAISON GROUP  
OFFICE OF NAVAL RESEARCH  
AMERICAN EMBASSY - TOKYO  
AFC SAN FRANCISCO 96503 1

PROF. J. A. MUCKSTADT  
DEPT. OF OPERATIONS RESEARCH  
CORNELL UNIVERSITY  
ITHACA,  
NEW YORK  
14850 1

DR. JANET M. MYRE  
THE INSTITUTE OF DECISION SCIENCE  
FOR BUSINESS AND PUBLIC POLICY  
CLAREMONT MEN'S COLLEGE  
CLAREMONT  
CA 91711 1

MR. H. NISSELSOHN  
BUREAU OF THE CENSUS  
ROOM 2025  
FEDERAL BUILDING 3  
WASHINGTON ,  
D. C. 20533 1

NO. OF COPIES

MISS B. S. ORLEANS  
NAVAL SEA SYSTEMS COMMAND  
(SEA 03F)  
RM 10S08  
ARLINGTON VIRGINIA 20360

1

PROF. C. B OWEN  
DEPARTMENT OF STATISTICS  
SOUTHERN METHODIST UNIVERSITY  
DALLAS  
TEXAS  
75222

1

DR. A. PETRASOVITS  
ROOM 207B , FOOD AND DRUG ELDG.  
TUNNEY'S PASTURE  
OTTAWA , CANADIA K1A-0L2 ,  
CANADA

1

PROF. S. L. PFCENIX  
SIBLEY SCHOOL OF MECHANICAL AND  
AEROSPACE ENGINEERING  
CORNELL UNIVERSITY  
ITHACA  
NY

1

14850

DR. A. L. POWELL  
DIRECTOR  
OFFICE OF NAVAL RESEARCH BRANCH OFF  
495 SUMMER STREET  
BOSTON  
MA

1

02210

MR. F. R. PRIORI  
CCDE 224 OPERATIONS TEST AND ONRS  
EVALUATION FORCE (OPTEVFOR)  
NORFOLK ,  
VIRGINIA  
20360

1

PROF. M. L. PURI  
DEPT. OF MATHEMATICS  
P.O. BOX F  
INDIANA UNIVERSITY FOUNDATION  
BLOOMINGTON  
IN

1

47401

PROF. H ROBBINS  
DEPARTMENT OF MATHEMATICS  
COLUMBIA UNIVERSITY  
NEW YORK,  
NEW YORK 10027

1

PROF. M ROSENBLATT  
DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF CALIFORNIA SAN DIEGO  
LA JOLLA  
CALIFORNIA

1

92093

PROF. S. M. ROSS  
COLLEGE OF ENGINEERING  
UNIVERSITY OF CALIFORNIA  
BERKELEY  
CA

1

94720

PROF. I RUBIN  
SCHOOL OF ENGINEERING AND APPLIED  
SCIENCE  
UNIVERSITY OF CALIFORNIA  
LOS ANGELES  
CALIFORNIA 90024

1

PROF. J. R. SAVAGE  
DEPARTMENT OF STATISTICS  
YALE UNIVERSITY  
NEW HAVEN,  
CONNECTICUT  
06520

1

PROF. L. L. , JR. SCHARF  
DEPARTMENT OF ELECTRICAL ENGINEERING  
COLORADO STATE UNIVERSITY  
FT. COLLINS ,  
COLORADO  
80521

1

PROF. R. SERFLING  
DEPARTMENT OF STATISTICS  
FLORIDA STATE UNIVERSITY

1

TALLAHASSEE FLORIDA 32306

PROF. W. R. SCHUCANY  
DEPARTMENT OF STATISTICS  
SOUTHERN METHODIST UNIVERSITY  
DALLAS ,  
TEXAS  
75222

1

PROF. D. O. SIEGMUND  
DEPT. OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD  
CA

1

94305

PROF. M. L. SHOUMAN  
DEPT. OF ELECTRICAL ENGINEERING  
POLYTECHNIC INSTITUTE OF NEW YORK  
BROOKLYN,  
NEW YORK  
11201

1

PROF. N. SINGPURWALLA  
DEPT. OF OPERATIONS RESEARCH  
THE GEORGE WASHINGTON UNIVERSITY  
707 22ND ST. N. W.  
WASHINGTON, D. C.

1

31 20052



NO. OF COPIES

DR. A. L. SLAFKOSKY 1  
SCIENTIFIC ADVISOR  
COMMANCANT OF THE MARINE CORPS  
WASHINGTON ,  
C. C.  
20380

MR. CHARLES S. SMITH 1  
OASD (I&L),  
PENTAGON  
WASHINGTON  
DC

20301

DR. D. E. SMITH 1  
DESMATICS INC.  
P.O. BOX 618  
STATE COLLEGE  
PENNSYLVANIA  
16801

PROF. W. L. SMITH 1  
DEPARTMENT OF STATISTICS  
UNIVERSITY OF NORTH CAROLINA  
CHAPEL HILL  
NORTH CAROLINA 27514

PROF. H. SOLOMON 1  
DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD ,  
CALIFORNIA  
94305

MR. GLENN F. STAHLY 1  
NATIONAL SECURITY AGENCY  
FORT MEADE ,  
MARYLAND 20755

MR. DAVID A. SWICK 1  
ADVANCED PROJECTS GROUP  
CODE 8103  
NAVAL RESEARCH LAB.  
WASHINGTON  
DC

20375

MR. WENDELL G. SYKES 1  
ARTHUR D. LITTLE, INC.  
ACORN PARK  
CAMBRIDGE  
MA

02140

PROF. J. R. THOMPSON 1  
DEPARTMENT OF MATHEMATICAL SCIENCE  
RICE UNIVERSITY  
HOUSTON,  
TEXAS  
77001



PROF. W. A. THOMPSON  
DEPARTMENT OF STATISTICS  
UNIVERSITY OF MISSOURI  
COLUMBIA,  
MISSOURI  
65201

1

PROF. F. A. TILLMAN  
DEPT. OF INDUSTRIAL ENGINEERING  
KANSAS STATE UNIVERSITY  
MANHATTAN  
KS

1

66506

PROF. J. W. TUKEY  
DEPARTMENT OF STATISTICS  
PRINCETON UNIVERSITY

1

PRINCETON, N. J. 08540

PROF. A. F. VEINOTT  
DEPARTMENT OF OPERATIONS RESEARCH  
STANFORD UNIVERSITY  
STANFORD  
CALIFORNIA  
94305

1

DANIEL H. WAGNER  
STATION SQUARE ONE  
EAGLE, PENNSYLVANIA  
19301

1

PROF. GRACE WAHBA  
DEPT. OF STATISTICS  
UNIVERSITY OF WISCONSIN  
MADISON  
WI

1

53706

PROF. K. T. WALLENIUS  
DEPARTMENT OF MATHEMATICAL SCIENCES  
CLEMSON UNIVERSITY  
CLEMSON,  
SOUTH CAROLINA 29631

1

PROF. G. S. WATSON  
DEPARTMENT OF STATISTICS  
PRINCETON, N. J. 08540

1

PROF. BERNARD WIDROW  
STANFORD ELECTRONICS LAB  
STANFORD UNIVERSITY  
STANFORD  
CA

1

94305

PROF. G. E. WHITEHOUSE  
DEPT. OF INDUSTRIAL ENGINEERING  
LEHIGH UNIVERSITY  
BETHLEHEM  
PA

1

18015

NO. OF COPIES

PROF. S. ZACKS 1  
DEPT. OF MATHEMATICS AND STATISTICS  
CASE WESTERN RESERVE UNIVERSITY  
CLEVELAND.,  
CHIC  
44106

PROF. M. ZIA-FASSAN 1  
DEPARTMENT OF INDUSTRIAL AND  
SYSTEMS ENGINEERING  
ILLINOIS INSTITUTE OF TECHNOLOGY  
CHICAGO 60616  
IL

DR. DAVID BRILLINGER 1  
STATISTICS DEPT.  
UNIVERSITY OF CALIFORNIA  
BERKELEY,  
CALIFORNIA  
94720

DR. CHIN CHEN 1  
COMMONWEALTH ASSOCIATES, INC.  
209 E. WASHINGTON  
JACKSON, MI  
49201

PROFESSOR PETER CHEN 1  
ALFRED P. SLICAN SCHOOL OF MGMT  
MASS INSTITUTE OF TECHNOLOGY  
50 MEMORIAL DRIVE  
CAMBRIDGE,  
MASSACHUSETTS 02139

DR. E. CINLAR 1  
O.R. DEPT.  
NORTHWESTERN UNIVERSITY  
EVANSTON,  
ILLINOIS  
60201

DR. GUY C. CORYNEN 1  
L. 156 UNIV OF CALIFORNIA  
LAWRENCE LIVERMORE LABORATORY  
P. O. BOX 808  
LIVERMORE, CALIFORNIA  
94550

DR. MICHAEL A. CRANE 1  
CONTROL ANALYSIS CORPORATION  
800 WELCH ROAD  
PALO ALTO, CALIFORNIA 94304

DR. D. F. CALEY 1  
STATISTICS DEPT. (I A S)  
AUSTRALIAN NATIONAL UNIVERSITY  
CANBERRA, A. C. T. 2606  
AUSTRALIA

NO. OF COPIES

HSU DER-ANN 1  
SCH OF ENGR./APPLIED SCI.  
DEPT. OF CIVIL & GEOLOGICAL ENGR.  
THE ENGINEERING QUADRANGLE  
PRINCETON  
NEW JERSEY 08540

ASSISTANT DEAN DAV ELDREDGE 1  
SCHOOL OF BUSINESS ADMINISTRATION  
UNIVERSITY OF EVANSVILLE  
EVANSVILLE  
INDIANA  
47702

DR. G. W. FISHMAN 1  
CURRICULUM IN O. R.  
UNIVERSITY OF NORTH CAROLINA  
PHILLIPS ANNEX , CHAPPEL HILL  
NORTH CAROLINA  
27514

DR. A. V. HOLLEN 1  
DEPARTMENT OF PHYSIOLOGY  
THE UNIVERSITY  
LEEDS LS2 9JT  
ENGLAND

DR. EUGENE D. HOMER 1  
MANAGEMENT SCI. & ENGINEERING DEPT.  
C.W. POST COLLEGE  
OF LONG ISLAND UNIVERSITY  
GREENVALE, NEW YORK  
11548

DR. D. VERE JONES 1  
DEPT. OF MATHEMATICS ,  
VICTORIA UNIVERSITY OF WELLINGTON ,  
P. O. BOX 196  
WELLINGTON , NEW ZEALAND

J. M. LIITTSCHWAGER 1  
SYSTEMS DIVISION  
COLLEGE OF ENGINEERING  
THE UNIVERSITY OF IOWA  
IOWA CITY , IOWA  
52242

Y. W. DE KWAADSTENIET 1  
DEPARTMENT OF BIOLOGY  
FREE UNIVERSITY  
DE BOELELAAN 1087  
AMSTERDAM-PUITENVELDER  
THE NETHERLANDS

JACK P. LANCOLT 1  
DEFENCE & CIVIL INST. OF ENVIR. MED  
1133 SHEPARD AVE. WEST  
PO BOX 2000 DOWNSVIEW  
ONTARIO,  
CANADA

NO. OF COPIES

MR. FRANK MCNOLTY 1  
1142 PCME AVE.  
SUNNYVALE, CA  
94087

DR. P. A. F. MCNAN 1  
STATISTICS DEPT. (I A S)  
AUSTRALIAN NATIONAL UNIVERSITY  
CANBERRA, A. C. T 2606  
AUSTRALIA

G. W. MORRISON 1  
OAK RIDGE NATIONAL LABORATORY  
OAK RIDGE, TENNESSEE  
37830

MR. BRADLEY NICKY 1  
USCA. FOREST SERVICE  
P. O. BOX 5007  
FIVERSIDE  
CALIFORNIA  
92507

DR. DAVID CATES 1  
DEPT. OF STATISTICS  
HARVARD UNIVERSITY  
SCIENCE CENTER, 1 OXFORD ST.  
CAMBRIDGE, MA.  
02132

Professor Emanuel Parzen, Director 1  
Institute of Statistics  
Texas A&M University  
College Station, TX 77843

MR ART PETERSON 1  
DEPT. OF BIOSTATISTICS SC-32  
UNIVERSITY OF WASHINGTON  
SEATTLE, WASHINGTON  
98195

DAVID W. ROBINSON 1  
CONTROL ANALYSIS CORPORATION  
800 WELCH ROAD  
PALO ALTO  
CALIFORNIA  
94304

C. H. SAUER (21-129) 1  
THOMAS J. WATSON RESEARCH CENTER  
F.C. BOX 219  
YORKTOWN HEIGHTS, NEW YORK  
10598

NO. OF COPIES

MR. C. A. STEELE JR 1  
F.C. BOX 45  
MAGNOLIA, MA  
01930

PROF P. SULLO 1  
SCHOOL OF MANAGEMENT  
FENSSELAER POLYTECHNIC  
INSTITUTE  
TROY, N.Y.  
12181

DR. ROBERT TRAPPL 1  
INSTITUT FÜR ALLGEMEINE UND  
VERGLEICHENDE PHYSIOLOGIE DER  
UNIVERSITÄT WIEN  
SCHWARZSPANIERSTRASSE 17  
A-1090 WIEN ÖSTERREICH (AUSTRIA)

PALL A. WILLIS 1  
POLY TECHNICA  
2824 W. GEORGE MASON  
FALLS CHURCH, VA. 22042

TUNCAL M. YEGULALP 1  
KRLMB SCHOOL OF MINES  
918 S.W. MIDD  
COLUMBIA UNIVERSITY  
520 W. 120TH ST.  
NEW YORK, N.Y. 10027

SHIEGKI YOKOYAMA 1  
RESEARCH CENTER OF APPLIED  
INFORMATION SCIENCES  
TOHOKU UNIVERSITY  
KATAHIRA-CHO 2-1,  
SENDAI 980 JAPAN

Dean of Research 1  
Code 012  
Naval Postgraduate School  
Monterey, Ca. 93940

Library (Code 55) 1  
Naval Postgraduate School  
Monterey, CA 93940

Professor Peter Lewis 50  
Code 55Lw  
Naval Postgraduate School  
Monterey, Ca. 93940

R. J. Stampfel 1  
Code 55  
Naval Postgraduate School  
Monterey, Ca. 93940



U186305



JOSELYN KNOX LIBRARY - RESEARCH REPORTS

5 6853 01057927 9

~~U186305~~